

Whether you are just starting treatment or already taking JYNARQUE, the

ADPKD Peer Mentor Program and **ADPKD Peer Conversations Webinars** are here to support you every step of the way.

INDICATION:

JYNARQUE is a prescription medicine used to slow kidney function decline in adults who are at risk for rapidly progressing autosomal dominant polycystic kidney disease (ADPKD). It is not known if JYNARQUE is safe and effective in children.

SELECT IMPORTANT SAFETY INFORMATION:

Serious liver problems: JYNARQUE can cause serious liver problems that can lead to the need for a liver transplant or can lead to death. Stop taking JYNARQUE and call your healthcare provider right away if you get any of the following symptoms: feeling tired, loss of appetite, nausea, right upper stomach (abdomen) pain or tenderness, vomiting, fever, rash, itching, yellowing of the skin and white part of the eye (jaundice), dark urine.

Please read **FULL PRESCRIBING INFORMATION**, including **BOXED WARNING** and **MEDICATION GUIDE**.

Please read **IMPORTANT SAFETY INFORMATION** on pages 5-6.

PATIENT-LED

WEBINAR DISCUSSIONS

ADPKD Peer Conversations are interactive webinars covering a range of topics for patients and caregivers. Each webinar provides participants with the opportunity to connect with other members of the ADPKD community through Q&As and helpful discussions.

Topics discussed may include:

Strengthening your support system Talking about ADPKD with family Tips and tools for traveling



It is so important to meet other people with ADPKD, especially if you don't know anyone, because no one else quite understands what you're going through.

Diedre

Real ADPKD Patient



Hear Deidre's Story ▶

Click the QR code to register for an **ADPKD Peer Conversations Webinar**



Please read FULL PRESCRIBING INFORMATION, including BOXED WARNING and MEDICATION GUIDE.

ONE-ON-ONE **CONVERSATIONS**

WITH ADPKD PATIENTS

The ADPKD Patient Peer Mentor Program is an easy way to talk with someone else living with ADPKD who knows what you are going through. Connect with a Peer Mentor over a phone or video call to get answers to your ADPKD questions and get the support you've been looking for.

Topics discussed may include:

Navigating family conversations Talking to children about ADPKD Traveling while taking JYNARQUE Finding resources for support Becoming a patient advocate



Assembling your support team is key to managing ADPKD.

Fack

Real ADPKD Patient

Read Zack's Story ▶

Click the QR code to register for an ADPKD Peer Mentor



Please read FULL PRESCRIBING INFORMATION, including BOXED WARNING and MEDICATION GUIDE.

PERSONAL SUPPORT


when you need it

The ADPKD Patient Peer Mentor and ADPKD Peer Conversations Webinar Programs were designed specifically for you. They are great ways to connect with others who understand what you're going through.



It is very interesting to hear what people with rapidly progressing ADPKD have to say.

> Fernando Real ADPKD Patient







Click the QR code to hear from other patients

Patients were compensated for their time. Patient images reflect their health status at the time the photos and videos were taken.

Please read FULL PRESCRIBING INFORMATION, including BOXED WARNING and MEDICATION GUIDE.

INDICATION and IMPORTANT SAFETY INFORMATION for JYNARQUE® (tolvaptan)

INDICATION:

What is JYNARQUE?

JYNARQUE is a prescription medicine used to slow kidney function decline in adults who are at risk for rapidly progressing autosomal dominant polycystic kidney disease (ADPKD). It is not known if JYNARQUE is safe and effective in children.

IMPORTANT SAFETY INFORMATION:

- **Serious liver problems.** JYNARQUE can cause serious liver problems that can lead to the need for a liver transplant or can lead to death. Stop taking JYNARQUE and call your healthcare provider right away if you get any of the following symptoms:
 - feeling tired
 - loss of appetite
 - nausea
 - right upper stomach (abdomen) pain or tenderness
 - vomiting
 - fever

- rash
- itching
- yellowing of the skin and white part of the eye (jaundice)
- dark urine

It is important that you have a blood test before you start JYNARQUE to help reduce your risk of liver problems. Your healthcare provider will do a blood test to check your liver:

- before you start taking JYNARQUE
- at 2 weeks and 4 weeks after you start treatment with JYNARQUE
- then monthly for 18 months during treatment with JYNARQUE
- and every 3 months from then on

Because of the risk of serious liver problems, JYNARQUE is only available through a restricted distribution program called the JYNARQUE Risk Evaluation and Mitigation Strategy (REMS) Program.

Do not take JYNARQUE if you:

- have a history of liver problems or have signs or symptoms of liver problems, excluding polycystic liver disease
- cannot feel if you are thirsty or cannot replace fluids by drinking
- have been told that the amount of sodium (salt) in your blood is too high or too low
- · are dehydrated
- are allergic to tolvaptan or any of the ingredients in JYNARQUE
- · are unable to urinate

Tell your healthcare provider about all your medical conditions, including if you:

- have a history of sodium (salt) levels that are too low
- are pregnant or plan to become pregnant. It is not known
 if tolvaptan will harm your unborn baby. Tell your
 healthcare provider if you become pregnant or think
 that you may be pregnant
- are breastfeeding or plan to breastfeed. It is not known if tolvaptan passes into your breast milk. Do not breastfeed during your treatment with JYNARQUE. Talk to your healthcare provider about the best way to feed your baby during this time

IMPORTANT SAFETY INFORMATION for JYNARQUE® (tolvaptan) continued

Tell your healthcare provider about all the medicines you take, including prescription medicines, over-the-counter medicines, vitamins, and herbal supplements.

- Taking JYNARQUE with certain medicines could cause you to have too much tolvaptan in your blood. JYNARQUE should not be taken with certain medications. Your healthcare provider can tell you if it is safe to take JYNARQUE with other medicines
- Do not start taking a new medicine without talking to your healthcare provider

JYNARQUE may cause serious side effects, including:

- Too much sodium in your blood (hypernatremia) and loss of too much body fluid (dehydration). In some cases, dehydration can lead to extreme loss of body fluid called hypovolemia. You should drink water when you are thirsty and throughout the day and night. Stop taking JYNARQUE and call your healthcare provider if you cannot drink enough water for any reason, such as not having access to water, or vomiting or diarrhea. Tell your healthcare provider if you get any of the following symptoms:
 - dizziness
 - fainting
 - weight loss
 - a change in the way your heart beats
 - feeling confused or weak

What should you avoid while taking JYNARQUE?

Do not drink grapefruit juice during treatment with JYNARQUE. This could cause you to have too much tolvaptan in your blood.

The **most common side effects** of JYNARQUE are:

- thirst and increased fluid intake
- making large amounts of urine, urinating often, and urinating at night

These are not all the possible side effects of JYNARQUE. Talk to your healthcare provider about any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist.

If you have any questions about your health or medicines, talk to your healthcare professional.

To report SUSPECTED ADVERSE REACTIONS, contact Otsuka America Pharmaceutical, Inc. at 1-800-438-9927 or FDA at 1-800-FDA-1088 (www.fda.gov/medwatch).

Please read <u>FULL PRESCRIBING INFORMATION</u>, including **BOXED WARNING** and **MEDICATION GUIDE**.



